

Brooks Rand Labs Detection & Reporting Limits

Biota

Analyte	Method	MDL	MRL	Units
%TS	SM 2540G	0.30	1.00	%
Ag	EPA 1638	0.020	0.100	mg/kg
Al	EPA 1638	0.14	0.80	mg/kg
As	EPA 1638	0.11	0.30	mg/kg
As	EPA 1638 DRC	0.014	0.040	mg/kg
As(III)	EPA 1632	0.004	0.010	mg/kg
As(Inorg)	EPA 1632	0.004	0.010	mg/kg
As(Org)	EPA 1638	0.05	0.16	mg/kg
As(Org)	EPA 1638 DRC	0.014	0.040	mg/kg
As(V)	EPA 1632	0.004	0.010	mg/kg
B	EPA 1638	0.24	0.80	mg/kg
Ba	EPA 1638	0.03	0.16	mg/kg
Be	EPA 1638	0.030	0.090	mg/kg
Ca	EPA 1638	8.00	30.0	mg/kg
Cd	EPA 1638	0.007	0.020	mg/kg
Co	EPA 1638	0.020	0.160	mg/kg
Cr	EPA 1638	0.06	0.15	mg/kg
Cr	EPA 1638 DRC	0.018	0.120	mg/kg
Cu	EPA 1638	0.03	0.16	mg/kg
DMAs	EPA 1632	0.004	0.012	mg/kg
Fe	EPA 1638	1.2	5.0	mg/kg
Fe	EPA 1638 DRC	0.08	0.80	mg/kg
Hg	EPA 1631 Appendix	0.12	0.40	ng/g
Hg (low level)	EPA 1631 Appendix	0.04	0.10	ng/g
K	EPA 1638	1.5	4.5	mg/kg
Li	EPA 1638	0.025	0.120	mg/kg
MeHg	EPA 1630	1.00	3.00	ng/g
MeHg (low level)	EPA 1630	0.07	0.20	ng/g
Mg	EPA 1638	0.30	2.50	mg/kg
MMAAs	EPA 1632	0.004	0.012	mg/kg
Mn	EPA 1638	0.012	0.040	mg/kg
Mo	EPA 1638	0.010	0.030	mg/kg
Na	EPA 1638	7.0	20.0	mg/kg
Ni	EPA 1638	0.05	0.20	mg/kg
Ni	EPA 1638 DRC	0.05	0.20	mg/kg
Pb	EPA 1638	0.004	0.040	mg/kg
Sb	EPA 1638	0.007	0.020	mg/kg
Se	EPA 1638	0.12	0.30	mg/kg
Se	EPA 1638 DRC	0.06	0.15	mg/kg
Se(IV)	SOP BR-0061 HPLC	0.050	0.200	mg/kg
Se(VI)	SOP BR-0061 HPLC	0.050	0.200	mg/kg
Sn	EPA 1638	0.020	0.150	mg/kg
Sr	EPA 1638	0.008	0.070	mg/kg
Ti	EPA 1638	0.07	0.25	mg/kg

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.

Brooks Rand Labs Detection & Reporting Limits

Biota

Analyte	Method	MDL	MRL	Units
TI	EPA 1638	0.002	0.008	mg/kg
U	EPA 1638	0.008	0.025	mg/kg
V	EPA 1638	0.06	0.18	mg/kg
V	EPA 1638 DRC	0.014	0.120	mg/kg
Zn	EPA 1638	0.20	1.00	mg/kg

Sediment

Analyte	Method	MDL	MRL	Units
%TS	SM 2540G	0.30	1.00	%
Ag	EPA 1638	0.042	0.100	mg/kg
Al	EPA 1638	1.00	5.00	mg/kg
As	EPA 1632	0.010	0.030	mg/kg
As	EPA 1638	0.48	1.30	mg/kg
As	EPA 1638 DRC	0.05	0.36	mg/kg
As(III)	EPA 1632	0.040	0.120	mg/kg
As(Inorg)	EPA 1632	0.004	0.010	mg/kg
As(Org)	EPA 1638 DRC	0.05	0.36	mg/kg
As(V)	EPA 1632	0.040	0.120	mg/kg
Au	EPA 1638	0.05	0.15	mg/kg
B	EPA 1638	2.40	7.50	mg/kg
Ba	EPA 1638	0.11	1.00	mg/kg
Be	EPA 1638	0.24	0.72	mg/kg
Ca	EPA 1638	30.0	150	mg/kg
Cd	EPA 1638	0.023	0.070	mg/kg
Co	EPA 1638	0.06	0.50	mg/kg
Cr	EPA 1638	0.63	2.00	mg/kg
Cr	EPA 1638 DRC	0.16	1.00	mg/kg
Cu	EPA 1638	0.17	0.50	mg/kg
Cu	EPA 1638 DRC	0.49	2.50	mg/kg
DMAs	EPA 1632	0.005	0.015	mg/kg
Fe	EPA 1638	4.8	25.0	mg/kg
Fe	EPA 1638 DRC	4.2	25.0	mg/kg
Hg	EPA 1631 Appendix	0.15	0.50	ng/g
Hg (low level)	EPA 1631 Appendix	0.05	0.15	ng/g
Hg(ExtInorg)	EPA 3200	0.25	0.75	ng/g
Hg(ExtOrg)	EPA 3200	0.05	0.15	ng/g
Hg(F0)	SOP BR-0013	50.0	50.0	ng/g
Hg(F1)	SOP BR-0013	0.20	0.50	ng/g
Hg(F2)	SOP BR-0013	0.20	0.50	ng/g
Hg(F3)	SOP BR-0013	2.00	5.00	ng/g
Hg(F4)	SOP BR-0013	2.00	5.00	ng/g
Hg(F5)	SOP BR-0013	2.00	5.00	ng/g
Hg(F6)	SOP BR-0013	2.00	5.00	ng/g
Hg(II)R	SOP BR-0012	0.08	0.25	ng/g

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.

Brooks Rand Labs Detection & Reporting Limits

Sediment

<u>Analyte</u>	<u>Method</u>	<u>MDL</u>	<u>MRL</u>	<u>Units</u>
Hg(NonMob)	EPA 3200	0.25	0.75	ng/g
Hg(SemiMob)	EPA 3200	0.20	0.60	ng/g
K	EPA 1638	20	60	mg/kg
La	EPA 1638	0.005	0.050	mg/kg
MeHg	EPA 1630	0.008	0.025	ng/g
Mg	EPA 1638	2.5	25.0	mg/kg
MMAs	EPA 1632	0.004	0.012	mg/kg
Mn	EPA 1638	0.06	0.25	mg/kg
Mo	EPA 1638	0.10	1.00	mg/kg
Na	EPA 1638	8	75	mg/kg
Ni	EPA 1638	0.26	1.00	mg/kg
Ni	EPA 1638 DRC	0.23	1.00	mg/kg
Pb	EPA 1638	0.056	0.150	mg/kg
Sb	EPA 1638	0.05	0.15	mg/kg
Se	EPA 1638	0.90	1.80	mg/kg
Se	EPA 1638 DRC	0.16	1.00	mg/kg
Se(IV)	SOP BR-0061 HPLC	0.05	0.20	mg/kg
Se(VI)	SOP BR-0061 HPLC	0.05	0.20	mg/kg
Sn	EPA 1638	0.05	0.15	mg/kg
Sr	EPA 1638	0.020	0.060	mg/kg
Th	EPA 1638	0.25	0.75	mg/kg
Ti	EPA 1638	0.74	2.25	mg/kg
Tl	EPA 1638	0.018	0.050	mg/kg
U	EPA 1638	0.010	0.050	mg/kg
V	EPA 1638	5.20	15.0	mg/kg
V	EPA 1638 DRC	0.08	0.75	mg/kg
Zn	EPA 1638	0.40	1.80	mg/kg
Zn	EPA 1638 DRC	0.25	1.00	mg/kg

Water (Saline)

<u>Analyte</u>	<u>Method</u>	<u>MDL</u>	<u>MRL</u>	<u>Units</u>
Ag	EPA 1640 APDC	0.005	0.015	µg/L
Ag	EPA 1640 RP	0.011	0.036	µg/L
As	EPA 1640 RP	0.05	0.15	µg/L
Be	EPA 1640 RP	0.006	0.013	µg/L
Cd	EPA 1640 APDC	0.004	0.030	µg/L
Cd	EPA 1640 Column	0.0010	0.0100	µg/L
Cd	EPA 1640 RP	0.004	0.012	µg/L
Co	EPA 1640 Column	0.0020	0.0100	µg/L
Co	EPA 1640 RP	0.020	0.090	µg/L
Cr	EPA 1640 RP	0.040	0.100	µg/L
Cu	EPA 1640 APDC	0.09	0.55	µg/L
Cu	EPA 1640 Column	0.0040	0.0200	µg/L
Cu	EPA 1640 RP	0.04	0.12	µg/L

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.

Brooks Rand Labs Detection & Reporting Limits

Water (Saline)

Analyte	Method	MDL	MRL	Units
Fe	EPA 1640 Column	0.04	0.12	µg/L
Mn	EPA 1640 Column	0.00	0.00	µg/L
Ni	EPA 1640 APDC	0.040	0.120	µg/L
Ni	EPA 1640 Column	0.0060	0.0200	µg/L
Ni	EPA 1640 RP	0.04	0.14	µg/L
Pb	EPA 1640 APDC	0.015	0.150	µg/L
Pb	EPA 1640 Column	0.0010	0.0100	µg/L
Pb	EPA 1640 RP	0.003	0.013	µg/L
Sb	EPA 1640 RP	0.004	0.015	µg/L
Se	EPA 1640 RP	0.070	0.210	µg/L
Tl	EPA 1640 RP	0.003	0.010	µg/L
V	EPA 1640 Column	0.007	0.025	µg/L
V	EPA 1640 RP	0.015	0.100	µg/L
Zn	EPA 1640 Column	0.015	0.050	µg/L
Zn	EPA 1640 RP	0.26	0.75	µg/L

Water *

Analyte	Method	MDL	MRL	Units
Ag	EPA 1638	0.005	0.020	µg/L
Al	EPA 1638	0.50	1.50	µg/L
As	EPA 1632	0.030	0.100	µg/L
As	EPA 1638	0.09	0.27	µg/L
As	EPA 1638 DRC	0.009	0.030	µg/L
As(III)	EPA 1632	0.008	0.025	µg/L
As(III)	SOP BR-0061 HPLC	0.013	0.050	µg/L
As(Inorg)	EPA 1632	0.008	0.025	µg/L
As(V)	EPA 1632	0.008	0.025	µg/L
As(V)	SOP BR-0061 HPLC	0.013	0.050	µg/L
Au	EPA 1638	0.006	0.020	µg/L
B	EPA 1638	0.40	2.00	µg/L
Ba	EPA 1638	0.010	0.050	µg/L
Be	EPA 1638	0.050	0.150	µg/L
Bi	EPA 1638	0.010	0.050	µg/L
Ca	EPA 1638	6.00	30.0	µg/L
Cd	EPA 1638	0.007	0.020	µg/L
Cd	EPA 1638 DRC	0.002	0.010	µg/L
Ce	EPA 1638	0.0010	0.0100	µg/L
Co	EPA 1638	0.020	0.100	µg/L
Co	EPA 1638 DRC	0.010	0.100	µg/L
Cr	EPA 1638	0.05	0.15	µg/L
Cr	EPA 1638 DRC	0.015	0.045	µg/L
Cr(III)	EPA 1638	4.00	10.0	µg/L
Cr(VI)	SM3500-CrB/7196A mod.	4.0	10.0	µg/L
Cr(VI)	SOP BR-0061 HPLC	0.060	0.200	µg/L

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.

Brooks Rand Labs Detection & Reporting Limits

Water *

Analyte	Method	MDL	MRL	Units
Cu	EPA 1638	0.040	0.120	µg/L
Cu	EPA 1638 DRC	0.011	0.050	µg/L
DMAAs	EPA 1632	0.018	0.050	µg/L
DMAAs	SOP BR-0061 HPLC	0.013	0.050	µg/L
Fe	EPA 1638	1.4	5.0	µg/L
Fe	EPA 1638 DRC	0.24	1.00	µg/L
Fe	SM 3500-Fe B mod.	7.3	20.0	µg/L
Fe	SM 3500-Fe B mod.	7.3	20.0	µg/L
Fe(II)	SM 3500-Fe B mod.	7.8	20.0	µg/L
Fe(III)	SM 3500-Fe B mod.	7.8	20.0	µg/L
Hardness	EPA 1638	6.00	30.0	µg/L
Hg	EPA 1631	0.15	0.40	ng/L
Hg(AcidLab)	SOP BR-0003	0.15	0.40	ng/L
Hg(II)R	In-House	0.15	0.40	ng/L
Hg(TotVol)	SOP BR-0005	0.10	0.25	ng/L
In	EPA 1638	0.0020	0.0040	µg/L
K	EPA 1638	5.0	15.0	µg/L
La	EPA 1638	0.0010	0.0100	µg/L
Li	EPA 1638	0.05	0.15	µg/L
Lu	EPA 1638	0.0010	0.0100	µg/L
MeHg	EPA 1630	0.020	0.050	ng/L
MeHg (low level)	EPA 1630	0.010	0.025	ng/L
Mg	EPA 1638	3.00	15.0	µg/L
MMAAs	EPA 1632	0.012	0.050	µg/L
MMAAs	SOP BR-0061 HPLC	0.013	0.050	µg/L
Mn	EPA 1638	0.020	0.060	µg/L
Mn(II)	EPA 1638	0.020	0.060	µg/L
Mn(IV)	EPA 1638	0.020	0.060	µg/L
Mo	EPA 1638	0.010	0.030	µg/L
Na	EPA 1638	3.0	15.0	µg/L
Ni	EPA 1638	0.04	0.20	µg/L
Ni	EPA 1638 DRC	0.050	0.200	µg/L
Pb	EPA 1638	0.006	0.025	µg/L
Sb	EPA 1638	0.010	0.040	µg/L
Se	EPA 1638	0.11	0.33	µg/L
Se	EPA 1638 DRC	0.020	0.060	µg/L
Se(IV)	SOP BR-0061 HPLC	0.030	0.100	µg/L
Se(VI)	SOP BR-0061 HPLC	0.030	0.100	µg/L
SeCN	SOP BR-0061 HPLC	0.03	0.10	µg/L
Si	EPA 1638	100	300	µg/L
Sn	EPA 1638	0.050	0.150	µg/L
Sr	EPA 1638	0.010	0.050	µg/L
Tb	EPA 1638	0.0010	0.0100	µg/L
Te	EPA 1638	0.020	0.060	µg/L

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.

Brooks Rand Labs Detection & Reporting Limits

Water *

<u>Analyte</u>	<u>Method</u>	<u>MDL</u>	<u>MRL</u>	<u>Units</u>
Th	EPA 1638	0.003	0.020	µg/L
Ti	EPA 1638	0.05	0.20	µg/L
Tl	EPA 1638	0.003	0.010	µg/L
Tm	EPA 1638	0.0010	0.0100	µg/L
TSS	EPA 160.2	0.3	1.0	mg/L
U	EPA 1638	0.003	0.010	µg/L
V	EPA 1638	0.05	0.15	µg/L
V	EPA 1638 DRC	0.010	0.030	µg/L
W	EPA 1638	0.010	0.040	µg/L
Y	EPA 1638	0.0010	0.0100	µg/L
Zn	EPA 1638	0.06	0.20	µg/L
Zn	EPA 1638 DRC	0.028	0.200	µg/L

Quoted detection limits may be subject to change based on necessary dilutions. Reporting limits for elements not listed will be based on the method blanks prepared with the batch and the low standard used in the calibration. All solid matrices are on a wet-weight basis.

* For ICP-MS analyses, Industrial waste water (i.e. FGD) will most always be diluted 50x, Blood samples will be diluted 50x, and Urine samples will be diluted 10x.